



Download Toppr - India's best learning app for classes 5th to 12th

360° learning with our adaptive platform



Online Classes

Learn for free with short videos and live classes



Adaptive Practice

Practice smart with questions created for your unique needs



Mock Tests

Be exam ready by solving all India tests and previous years' papers



Live Doubts

Chat with tutors and get your doubts resolved instantly, 24x7



Live Classes

Learn concepts and get tips from the best teachers with free Live Classes


Learn for free with the best videos




Polish your basics with fun games



Practice smart with personalised questions



Get your doubts cleared instantly, 24x7



Download the app for FREE now

GET A 5-DAY FREE TRIAL



9,184,321
HAPPY STUDENTS

492,461,127
QUESTIONS ATTEMPTED

A+
TESTS TAKEN

Doubts Answered

NCERT Solutions for Class 8 Subject-wise

- [Class 8 Maths](#)
- [Class 8 Science – Physics](#)
- [Class 8 Science – Biology](#)
- [Class 8 Science – Chemistry](#)
- [Class 8 Social Science – History](#)
- [Class 8 Geography](#)
- [Class 8 General Knowledge](#)
- [Class 8 Civics](#)

#463493

Topic: Operations of Polynomials

Identify the terms, their coefficients for each of the following expressions.

(i) $5xyz^2 - 3zy$

(ii) $1 + x + x^2$

(iii) $4x^2y^2 - 4x^2y^2z^2 + z^2$

(iv) $3 - pq + qr - rp$

(v) $\frac{x}{2} + \frac{y}{2} - xy$

(vi) $0.3a - 0.6ab + 0.5b$

Solution

Expression	Terms	Coefficients
$5xyz^2 - 3zy$	$5xyz^2$, $-3zy$	5 -3
$1 + x + x^2$	1, x , x^2	No coefficient, 1, 1
$4x^2y^2 - 4x^2y^2z^2 + z^2$	$4x^2y^2$, $-4x^2y^2z^2$, z^2	4, -4, 1
$3 - pqqr - rp$	3, $-pq$, $-rp$	No coefficient, -1, -1
	$\frac{x}{2}$	$\frac{1}{2}$
$\frac{x}{2} + \frac{y}{2} - xy$	$\frac{y}{2}$ $-xy$	$\frac{1}{2}$ -1
$0.3a - 0.6ab + 0.5b$	$0.3a$ $-0.6ab$ $0.5b$	0.3, -0.6, 0.5

#463673

Topic: Special Products

Use suitable identities to find the following products:

(i) $(x + 4)(x + 10)$

(ii) $(x + 8)(x - 10)$

(iii) $(3x + 4)(3x - 5)$

(iv) $\left(y^2 + \frac{3}{2}\right)\left(y^2 - \frac{3}{2}\right)$

(v) $(3 - 2x)(3 + 2x)$

Solution

We use identities

$$(x + a)(x + b) = x^2 + (a + b)x + ab \text{ and } x^2 - y^2 = (x + y)(x - y)$$

$$(i) (x + 4)(x + 10)$$

$$= x^2 + (4 + 10)x + (4)(10)$$

$$= x^2 + 14x + 40$$

$$(ii) (x + 8)(x - 10)$$

$$= x^2 + (8 - 10) + (8)(-10)$$

$$= x^2 - 2x - 80$$

$$(iii) (3x + 4)(3x + 5)$$

$$= (3x)^2 + (4 - 5)3x + (4)(-5)$$

$$= 9x^2 - 3x - 20$$

$$(iv) \left(y^2 + \frac{3}{2}\right)\left(y^2 - \frac{3}{2}\right)$$

$$= (y^2)^2 - \left(\frac{3}{2}\right)^2$$

$$= y^4 - \frac{9}{4}$$

$$(v) (3 - 2x)(3 + 2x)$$

$$= (3)^2 - (2x)^2 = 9 - 4x^2$$

#463772

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$4(x - 5) = 4x - 5$$

Solution

The statement is incorrect.

The correct statement is

$$4(x - 5) = 4x - 20$$

#463773

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$x(3x + 2) = 3x^2 + 2$$

Solution

The given statement is incorrect.

The correct statement is

$$x(3x + 2) = 3x^2 + 2x$$

#463774

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$2x + 3y = 5xy$$

Solution

The given statement is incorrect.

As we cannot apply any rule over here, the correct statement will be

$$2x + 3y = 2x + 3y$$

#463775

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$x + 2x + 3x = 5x$$

Solution

The given statement is incorrect.

As all the terms have same variable x , we can add them.

Therefore, correct statement is

$$x + 2x + 3x = 6x$$

#463776

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$5y + 2y + y - 7y = 0$$

Solution

The given statement is incorrect.

As all the terms have same variable, we can perform operation of addition and subtraction.

Therefore, the correct statement is

$$5y + 2y + y - 7y = y$$

#463777

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$3x + 2x = 5x^2$$

Solution

The statement is incorrect.

As all the terms have same variable x , we can perform the operation.

Therefore the correct statement is

$$3x + 2x = 5x$$

#463778

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$(2x)^2 + 4(2x) + 7 = 2x^2 + 8x + 7$$

Solution

The given statement is incorrect.

Only the first term is written incorrect.

Correct statement is

$$(2x)^2 + 4(2x) + 7 = 4x^2 + 8x + 7$$

#463779

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$(2x)^2 + 5x = 4x + 5x = 9x$$

Solution

The given statement is incorrect.

Correct statement is

$$(2x)^2 + 5x = 4x^2 + 5x$$

#463780

Topic: Standard Identities

Find and correct errors of the following mathematical expressions:ind and correct error

$$(3x + 2)^2 = 3x^2 + 6x + 4$$

Solution

The statement is incorrect.

LHS can be expanded by using formula, $(a + b)^2 = a^2 + 2ab + b^2$

$$\therefore (3x + 2)^2 = (3x)^2 + 2(3x)(2) + 2^2 = 9x^2 + 12x + 4$$

Therefore the correct statement is

$$(3x + 2)^2 = 9x^2 + 12x + 4$$

#463781

Topic: Algebraic Expressions and Types

Find and correct error in the following mathematical statement.

Substituting $x = -3$ in

$$(a) x^2 + 5x + 4 \text{ gives } (-3)^2 + 5(-3) + 4 = 9 + 2 + 4 = 15$$

$$(b) x^2 - 5x + 4 \text{ gives } (-3)^2 - 5(-3) + 4 = 9 - 15 + 4 = -2$$

$$(c) x^2 + 5x \text{ gives } (-3)^2 + 5(-3) = -9 - 15 = -24$$

Solution

$$(a) x^2 + 5x + 4 = (-3)^2 + 5(-3) + 4 = 9 - 15 + 4 = -2$$

$$(b) x^2 - 5x + 4 = (-3)^2 - 5(-3) + 4 = 9 + 15 + 4 = 28$$

$$(c) x^2 + 5x = (-3)^2 + 5(-3) = 9 - 15 = -6$$

#463782

Topic: Standard Identities

Find and correct errors of the following mathematical expressions:

$$(y - 3)^2 = y^2 - 9$$

Solution

The statement is incorrect.

It can be expanded by using formula $(a - b)^2$. But instead of this the formula used here is $a^2 - b^2$ which is incorrect.

Therefore the correct statement is

$$(y - 3)^2 = y^2 - 2 \times y \times 3 + 9 = y^2 - 6y + 9$$

#463783

Topic: Standard Identities

Find and correct errors of the following mathematical expressions:

$$(z + 5)^2 = z^2 + 25$$

Solution

The statement is incorrect.

Above expression can be expanded by using formula $(a + b)^2 = a^2 + 2ab + b^2$

The correct statement is

$$(z + 5)^2 = z^2 + 10z + 25$$

#463784

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$(a + 4)(a + 2) = a^2 + 8$$

Solution

The given statement is incorrect.

Correct statement is

$$(a + 4)(a + 2) = a^2 + 6a + 8$$

#463785

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$(a - 4)(a - 2) = a^2 - 8$$

Solution

The given statement is incorrect. Middle term is missing in this expression.

Correct statement is

$$(a - 4)(a - 2) = a^2 - 6a + 8$$

#463786

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$(2a + 3b)(a - b) = 2a^2 - 3b^2$$

Solution

The given statement is incorrect. Middle term is missing in the expression.

Correct statement is

$$(2a + 3b)(a - b) = 2a^2 + ab - 3b^2$$

#463787

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$\frac{3x^2}{3x^2} = 0$$

Solution

The statement is incorrect. Like terms after cancellation leave remainder as 1.

Therefore the correct statement is

$$\frac{3x^2}{3x^2} = 1$$

#463788

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$\frac{3x^2 + 1}{3x^2} = 1 + 1 = 2$$

Solution

The given statement is incorrect.

After separating the denominators, we get

$$\frac{3x^2 + 1}{3x^2} = 1 + \frac{1}{3x^2}$$

The above statement is the correct one.

#463789

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$\frac{3x}{3x + 2} = \frac{1}{2}$$

Solution

The given statement is incorrect.

As we cannot separate the denominators, the statement will remain as it is.

Therefore correct statement is

$$\frac{3x}{3x + 2} = \frac{3x}{3x + 2}$$

#463790

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$\frac{3}{4x + 3} = \frac{1}{4x}$$

Solution

The given statement is incorrect.

As we cannot separate the denominators, the statement will remain as it is.

Therefore, correct statement is

$$\frac{3}{4x + 3} = \frac{3}{4x + 3}$$

#463791

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$\frac{4x + 5}{4x} = 5$$

Solution

The given statement is incorrect.

After separating the denominators, we get

$$\frac{4x + 5}{4x} = 1 + \frac{5}{4x} \text{ which is the correct statement.}$$

#463792

Topic: Operations of Polynomials

Find and correct errors of the following mathematical expressions:

$$\frac{7x + 5}{5} = 7x$$

Solution

The given statement is incorrect.

We will get correct statement by separating denominators, which is

$$\frac{7x+5}{5} = 1 + \frac{7x}{5}$$

#463838

Topic: Standard Form of Polynomial

Classify the following polynomials as monomials, binomials, and trinomials. Which polynomial do not fit in any of these three categories?

$x+y$, 1000 , $x+x^2+x^3+x^4$, $7+y+5x$, $2y-3y^2$, $2y-3y^2+4y^3$, $5x-4y+3xy$, $4z-15z^2$, $ab+bc+cd+da$, pqr , p^2q+pq^2 , $2p+2q$

Solution

Monomials :

1000 , pqr

Binomials :

$x+y$, $2y-3y^2$, $4z-15z^2$, p^2q+pq^2 , $2p+2q$

Trinomials :

$7+y+5x$, $2y-y^2+4y^3$, $5x-4y+3xy$

Polynomials that do not fit in any of these categories are :

$x+x^2+x^3+x^4$, $ab+bc+cd+da$

#463862

Topic: Operations of Polynomials

Carry out the following divisions

(i) $28x^4 \div 56x$

(ii) $-36y^3 \div 9y^2$

(iii) $66pq^2r^3 \div 11qr^2$

(iv) $34x^3y^3z^3 \div 51xy^2z^3$

(v) $12a^8b^8 \div (-6a^6b^4)$

Solution

(i) $28x^4 \div 56x$

$$= \frac{28x^4}{56x} = \frac{x^3}{2}$$

(ii) $-36y^3 \div 9y^2$

$$= \frac{-36y^3}{9y^2} = -4y$$

(iii) $66pq^2r^3 \div 11qr^2$

$$= \frac{66pq^2r^3}{11qr^2} = 6pqr$$

(iv) $34x^3y^3z^3 \div 51xy^2z^3$

$$= \frac{34x^3y^3z^3}{51xy^2z^3} = \frac{2}{3}x^2y$$

(v) $12a^8b^8 \div (-6a^6b^4)$

$$\frac{12a^8b^8}{(-6a^6b^4)} = -2a^2b^4$$

#463881

Topic: Operations of Polynomials

Divide the polynomial by the given monomial

(i) $(5x^2 - 6x) \div 3x$

(ii) $(3y^8 - 4y^6 + 5y^4) \div y^4$

(iii) $8(x^3y^2z^2 + x^2y^3z^2 + x^2y^2z^3) \div 4x^2y^2z^2$

(iv) $(x^3 + 2x^2 + 3x) \div 2x$

(v) $(p^3q^6 - p^6q^3) \div p^3q^3$

Solution

(i) $(5x^2 - 6x) \div 3x$

= \frac{1}{3}(5x - 6)

(ii) $(3y^8 - 4y^6 + 5y^4) \div y^4$

$$= 3y^4 - 4y^2 + 5$$

(iii) $8(x^3y^2z^2 + x^2y^3z^2 + x^2y^2z^3) \div 4x^2y^2z^2$

$$= 2(x + y + z)$$

(iv) $(x^3 + 2x^2 + 3x) \div 2x$

$$= \frac{1}{2}(x^2 + 2x + 3)$$

(v) $(p^3q^6 - p^6q^3) \div p^3q^3$

$$= q^3 - p^3$$

#463887

Topic: Operations of Polynomials

Work out the following division

(i) $(10x - 25) \div 5$

(ii) $(10x - 25) \div (2x - 5)$

(iii) $10y(6y + 21) \div 5(2y + 7)$

(iv) $9x^2y^2(3z - 24) \div 27xy(z - 8)$

(v) $96abc(3a - 12)(5b - 30) \div 144(a - 4)(b - 6)$

Solution

(i) $(10x - 25) \div 5 = 2x - 5$

(ii) $(10x - 25) \div (2x - 5) = 5$

(iii) $10y(6y + 21) \div 5(2y + 7) = 6y$

(iv) $9x^2y^2(3z - 24) \div 27xy(z - 8) = xy$

(v) $96abc(3a - 12)(5b - 30) \div 144(a - 4)(b - 6) = 10abc$

#463896

Topic: Operations of Polynomials

Divide as directed

- (i) $5(2x + 1)(3x + 5) \div (2x + 1)$
- (ii) $26xy(x + 5)(y - 4) \div 13x(y - 4)$
- (iii) $52pqr(p + q)(q + r)(r + p) \div 104pq(q + r)(r + p)$
- (iv) $20(y + 4)(y^2 + 5y + 3) \div 5y(y + 4)$
- (v) $x(x + 1)(x + 2)(x + 3) \div x(x + 1)$

Solution

(i) $5(2x + 1)(3x + 5) \div (2x + 1) = 5(3x + 1)$

(ii) $26xy(x + 5)(y - 4) \div 13x(y - 4) = 2y(x + 5)$

(iii) $52pqr(p + q)(q + r)(r + p) \div 104pq(q + r)(r + p) = \frac{1}{2}r(p + q)$

(iv) $20(y + 4)(y^2 + 5y + 3) \div 5y(y + 4) = 4(y^2 + 5y + 3)$

(v) $x(x + 1)(x + 2)(x + 3) \div x(x + 1) = (x + 2)(x + 3)$

#463910

Topic: Operations of Polynomials

Add following.

- (i) $ab - bc, bc - ca, ca - ab$
- (ii) $a - b + ab, b - c + bc, c - a + ac$
- (iii) $2p^2q^2 - 3pq + 4, 5 + 7pq - 3p^2q^2$
- (iv) $l^2 + m^2, m^2 + n^2, n^2 + l^2, 2lm + 2mn + 2nl$

Solution

i) $ab - bc + bc - ca + ca - ab = 0$

ii) $a - b + ab + b - c + bc + c - a + ac = ab + bc + ac$

iii) $2p^2q^2 - 3pq + 4 + 5 + 7pq - 3p^2q^2 = -p^2q^2 + 4pq + 9$

iv) $l^2 + m^2 + n^2 + m^2 + l^2 + n^2 + 2lm + 2mn + 2nl = 2l^2 + 2m^2 + 2n^2 + 2(lm + mn + nl) = 2(l^2 + m^2 + n^2 + lm + mn + nl)$

#464017

Topic: Operations of Polynomials

Factorise the expression and divide them as directed

- (i) $(y^2 + 7y + 10) \div (y + 5)$
- (ii) $(m^2 - 14m - 32) \div (m + 2)$
- (iii) $(5p^2 - 25p + 20) \div (p - 1)$
- (iv) $4yz(z^2 + 6z - 16) \div 2y(z + 8)$
- (v) $(5pq(p^2 - q^2)) \div 2p(p + q)$
- (vi) $12xy(9x^2 - 16y^2) \div 4xy(3x + 4y)$
- (vii) $39y^3(50y^2 - 98) \div 26y^2(5y + 7)$

Solution

(i) $(y^2 + 7y + 10) \div (y + 5) = \frac{(y^2 + 5y + 2y + 10)}{y + 5}$

$= \frac{y(y + 5) + 2(y + 5)}{y + 5} = \frac{(y + 5)(y + 2)}{(y + 5)}$

$$= y + 2$$

$$(ii) (m^2 - 14m - 32) \div (m + 2) = \frac{(m^2 - 16m + 2m - 32)}{m + 2}$$

$$= \frac{m(m - 16) + 2(m - 16)}{m + 2} = \frac{(m - 16)(m + 2)}{(m + 2)}$$

$$= m - 16$$

$$(iii) (5p^2 - 25p + 20) \div (p - 1) = \frac{(5p^2 - 5p - 20p + 20)}{p - 1}$$

$$= \frac{5p(p - 1) - 20(p - 1)}{p - 1} = \frac{(p - 1)(5p - 20)}{(p - 1)}$$

$$= 5p - 20 = 5(p - 4)$$

$$(iv) 4yz(z^2 + 6z - 16) \div 2y(z + 8) = 4yz \frac{(z^2 + 8z - 2z - 16)}{2y(z + 8)}$$

$$= 2z \frac{z(z + 8) - 2(z + 8)}{z + 8} = 2z \frac{(z - 2)(z + 8)}{(z + 8)}$$

$$= 2z(z - 2)$$

$$(v) 5pq(p^2 - q^2) \div 2p(p + q)$$

$$= 5pq \frac{(p - q)(p + q)}{2p(p + q)}$$

$$= \frac{5}{2}q(p - q)$$

$$(vi) 12xy(9x^2 - 16y^2) \div 4xy(3x + 4y) = 12xy \frac{[(3x)^2 - (4y)^2]}{4xy(3x + 4y)}$$

$$= 3 \frac{(3x + 4y)(3x - 4y)}{(3x + 4y)}$$

$$= 3(3x - 4y)$$

$$(vii) 39y^3(50y^2 - 98) \div 26y^2(5y + 7) = 39y^3 \times 2 \frac{(25y^2 - 49)}{26y^2(5y + 7)}$$

$$= 3y \frac{[(5y)^2 - 7^2]}{5y + 7} = 3y \frac{[(5y - 7)(5y + 7)]}{5y + 7}$$

$$= 3y(5y - 7)$$

#464107

Topic: Operations of Polynomials

Subtract

- (i) $4a - 7ab + 3b + 12$ from $12a - 9ab + 5b - 3$
(ii) $3xy + 5yz - 7zx$ from $5xy - 2yz - 2zx + 10xyz$
(iii) $4p^2q - 3pq + 5pq^2 - 8p + 7q - 10$ from $18 - 3p - 11q + 5pq - 2pq^2 + 5p^2q$

Solution

i) $12a - 9ab + 5b - 3 - (4a - 7ab + 3b + 12) = 12a - 9ab + 5b - 3 - 4a + 7ab - 3b - 12 = 8a + 2b - 2ab - 15$

ii) $5xy - 2yz - 2zx + 10xyz - (3xy + 5yz - 7zx) = 5xy - 2yz - 2zx + 10xyz - 3xy - 5yz + 7zx = 2xy - 7yz + 5xz$

iii)

$$18 - 3p - 11q + 5pq - 2pq^2 + 5p^2q - (4p^2q - 3pq + 5q^2p - 8p + 7q - 10) = 18 - 3p - 11q + 5pq - 2pq^2 + 5p^2q - 4p^2q + 3pq - 5q^2p + 8p - 7q + 10 = 28 + 5p - 18q + 8pq - 7pq^2 + p^2q$$

#464110

Topic: Operations of Polynomials

Find product of following pairs of monomials

- (i) $4, 7p$
(ii) $-4p, 7p$
(iii) $-4p, 7pq$
(iv) $4p^3, -3p$
(v) $4p, 0$

Solution

(i) $4 \times 7 \times p = 28p$

(ii) $-4p \times 7p = -28p^2$

(iii) $-4p \times 7pq = -28p^2q$

(iv) $4p^3 \times 3p = -12p^4$

(v) $4p \times 0 = 0$

#464113

Topic: Operations of Polynomials

Find areas of rectangles with following pairs of monomials as their length and breadth respectively.

- (i) (p, q)
(ii) $10m, 5n$
(iii) $20x^2, 5y^2$
(iv) $(4x, 3x^2)$
(v) $3mn, 4np$

Solution

Area of rectangle, $A = \text{Length} \times \text{Breadth}$

(i) $A = l \times b = p \times q = pq$

(ii) $A = l \times b = 5n \times 10m = 50mn$

(iii) $A = l \times b = 20x^2 \times 5y^2 = 100x^2y^2$

(iv) $A = l \times b = 4x \times 3x^2 = 12x^3$

(v) $A = l \times b = 3mn \times 4np = 12m_n^2p$

#464155

Topic: Operations of Polynomials

Complete the table of products

<u>1st monomial</u> → <u>2nd monomial</u> ↓	$2x$	$-5y$	$3x^2$	$-4xy$	$7x^2y$	$-9x^2y^2$
$2x$	$4x^2$
$-5y$	$-15x^2y$
$3x^2$
$-4xy$
$7x^2y$
$-9x^2y^2$

Solution

<u>1st monomial</u> → <u>2nd monomial</u> ↓	$2x$	$-5y$	$3x^2$	$-4xy$	$7x^2y$	$-9x^2y^2$
$2x$	$4x^2$	$-10xy$	$6x^3$	$-8x^2y$	$14x^3y$	$-18x^3y^2$
$-5y$	$-10xy$	$25y^2$	$-15x^2y$	$20xy^2$	$-35x^2y^2$	$45x^2y^3$
$3x^2$	$6x^3$	$-15x^2y$	$9x^4$	$-12x^3y$	$21x^4y$	$-27x^4y^2$
$-4xy$	$-8x^2y$	$20xy^2$	$-12x^3y$	$16x^2y^2$	$-28x^3y^2$	$36x^3y^3$
$7x^2y$	$14x^3y$	$-35x^2y^2$	$21x^4y$	$-28x^3y^2$	$49x^4y^2$	$-63x^4y^3$
$-9x^2y^2$	$-18x^3y^2$	$45x^2y^3$	$-27x^4y^2$	$36x^3y^3$	$-63x^4y^3$	$81x^4y^4$

#464163

Topic: Operations of Polynomials

Obtain the volume of rectangular boxes with following length, breadth and height respectively.

(i) $5a, 3a^2, 7a^4$

(ii) $2p, 4q, 8r$

(iii) $xy, 2x^2y, 2xy^2$

(iv) $a, 2b, 3c$

Solution

Volume = Length × Breadth × Height

$$V = l \times b \times h$$

$$(i) V = 5a \times 3a^2 \times 7a^4 = 105a^7$$

$$(ii) V = l \times b \times h = 2p \times 4q \times 8r = 64pqr$$

$$(iii) V = l \times b \times h$$

$$V = xy \times 2x^2y \times 2xy^2$$

$$= 4x^4y^4$$

$$(iv) V = l \times b \times h = a \times 2b \times 3c = 6abc$$

#464168

Topic: Operations of Polynomials

Obtain the product of

$$(i) xy, yz, zx$$

$$(ii) a, -a^2, a^3$$

$$(iii) 2, 4y, 8y^2, 16y^3$$

$$(iv) a, 2b, 3c, 6abc$$

$$(v) m, -mn, mnp$$

Solution

$$(i) xy \times yz \times zx = x^2y^2z^2$$

$$(ii) a \times -a^2 \times a^3 = -a^6$$

$$(iii) 2 \times 4y \times 8y^2 \times 16y^3 = 1024y^6$$

$$(iv) a \times 2b \times 3c \times 6abc = 36a^2b^2c^2$$

$$(v) m \times -mn \times mnp = -m^3n^2p$$

#464179

Topic: Operations of Polynomials

Carry out the multiplication of expressions in each of the following pairs.

$$(i) 4p, q + r$$

$$(ii) ab, a - b$$

$$(iii) a + b, 7a^2b^2$$

$$(iv) (a^2 - 9) \times (4a)$$

$$(v) pq + qr + rp, 0$$

Solution

$$\begin{aligned} \text{(i)} \quad & (4p) \times (q + r) \\ & = (4p \times q) + (4p \times r) \\ & = 4pq + 4pr \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad & ab \times (a - b) \\ & = (ab \times a) + (ab \times (-b)) \\ & = a^2b^2 - ab^2 \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad & (a + b) \times (7a^2b^2) \\ & = (a \times 7a^2b^2) + (b \times 7a^2b^2) \\ & = 7a^3b^2 + 7a^2b^3 \end{aligned}$$

$$\begin{aligned} \text{(iv)} \quad & (a^2 \times 4a) + (-9 \times 4a) \\ & = 4a^3 - 36a \end{aligned}$$

$$\begin{aligned} \text{(v)} \quad & (pq \times 0) + (qr \times 0) + (rp \times 0) \\ & = 0 + 0 + 0 \\ & = 0 \end{aligned}$$

#464188**Topic:** Operations of Polynomials

Complete the table.

First expression	Second expression	Product
(i) a	$b + c + d$
(ii) $x + y - 5$	$5xy$
(iii) p	$6p^2 - 7p + 5$
(iv) $4p^2q^2$	$p^2 - q^2$
(v) $a + b + c$	abc

Solution

(i) $a(b + c + d) = ab + ac + ad$

(ii) $(x + y - 5)5xy = 5x^2y + 5xy^2 - 25xy$

(iii) $p(6p^2 - 7p + 5) = 6p^3 - 7p^2 + 5p$

(iv) $4p^2q^2(p^2 - q^2) = 4p^4q^2 - 4p^2q^4$

(v) $(a + b + c)abc = a^2bc + ab^2c + abc^2$

#464194**Topic:** Operations of Polynomials

Find the product.

(i) $a^2 \times (2a^{22}) \times (4a^{26})$

(ii) $\left(\frac{2}{3}xy\right) \times \left(-\frac{9}{10}x^2y^2\right)$

(iii) $\left(\frac{-10}{3}pq^3\right) \times \left(\frac{6}{5}p^3q\right)$

(iv) $x \times x^2 \times x \times x^3 \times x^4$

Solution

(i) As all the terms are having same base, then we can add the powers.

$$\therefore 8a^{2+22+26} = 8a^{50}$$

(ii) $\left(\frac{2}{3} \times y\right) \left(-\frac{9}{10}x^2y^2\right) = \frac{-3}{5}x^3y^3$

(iii) $\left(\frac{-10}{3}pq^3\right) \times \left(\frac{6}{5}p^3q\right) = -4p^4q^4$

(iv) $x \times x^2 \times x \times x^3 \times x^4 = x^{2+3+4} = x^{10}$

#464198

Topic: Operations of Polynomials

(a) Simplify $3x(4x - 5) + 3$ and find its value for

(i) $x = 3$ (ii) $x = \frac{1}{2}$

(b) Simplify $a(a^2 + a + 1) + 5$ and find its value for

(i) $a = 0$ (ii) $a = 1$ (iii) $a = -1$

Solution

$$\begin{aligned} \text{(a) (i)} & 3x(4x - 5) + 3 \\ & = 3 \times 3(4 \times 3 - 5) + 3 \\ & = 9(7) + 3 = 66 \end{aligned}$$

$$\begin{aligned} \text{(ii)} & 3x(4x - 5) + 3 \\ & = 3 \times \frac{1}{2}(4 \times \frac{1}{2} - 5) + 3 \\ & = \frac{3}{2}(2 - 5) + 3 \\ & = \frac{-3}{2} \end{aligned}$$

$$\begin{aligned} \text{[b] (i)} & a(a^2 + a + 1) + 5 \\ & = 0 + 5 \\ & = 5 \end{aligned}$$

$$\begin{aligned} \text{(ii)} & a(a^2 + a + 1) + 5 \\ & = 1(3) + 5 \\ & = 8 \end{aligned}$$

$$\begin{aligned} \text{(iii)} & a(a^2 + a + 1) + 5 \\ & = 1(1 - 1 + 1) + 5 \\ & = 1 + 5 \\ & = 4 \end{aligned}$$

#464215

Topic: Operations of Polynomials

- (a) Add: $p(p - q), q(q - r) \& r(r - p)$
 (b) Add: $2x(z - x - y) \& 27(z - y - x)$
 (c) Subtract: $3(l - 4m + 5n)$ from $4(l - 10n - 3m + 2l)$
 (d) Subtract: $3a(a + b + c) - 2b(a - b + c)$ from $4c(-a + b + c)$

Solution

$$\text{a) } p(p - q) + q(q - r) + r(r - p) = p^2 - pq + q^2 - qr + r^2 - pr = p^2 + q^2 + r^2 - pq - qr - rp$$

$$\text{b) } 2x(z - x - y) + 27(z - y - x) = 2xz - 2x^2 - 2xy + 27z - 27y - 27x$$

$$\text{c) } 4(l - 10n - 3m + 2l) - 3(l - 4m + 5n) = 40ln - 12lm + 8l^2 - 3l^2 + 12lm - 15ln = 25ln + 5l^2$$

$$\text{d) } 4c(-a + b + c) - [3a(a + b + c) - 2b(a - b + c)] = -4ac + 4bc + 4c^2 - [3a^2 + 3ab + 3ac - 2ab + 2b^2 - 2bc] = -4ac + 4bc + 4c^2 - 3a^2 - ab - 3ac - 2b^2 + 2bc$$

#464477

Topic: Operations of Polynomials

Multiply the binomials

(i) $(2x + 5)$ and $(4x - 3)$

(ii) $(y - 8)$ and $(3y - 4)$

(iii) $(2.5l - 0.5m)$ and $(2.5l + 0.5m)$

(iv) $(a + 3b)$ and $(x + 5)$

(v) $(2pq + 3q^2)$ and $(3pq - 2q^2)$

(vi) $\left(\frac{3}{4}a^2 + 3b^2\right)$ and $\left(a^2 - \frac{2}{3}b^2\right)$

Solution

(i) $2x \times (4x - 3) + 5 \times (4x - 3)$

$$= 8x^2 - 6x + 20x - 15$$

$$= 8x^2 + 14x - 15$$

(ii) $y(3y - 4) - 8(3y - 4)$

$$= 3y^2 - 28y + 32$$

(iii) $2.5l(2.5l + 0.5m) - 0.5m(2.5l + 0.5m)$

$$= 6.25l^2 - 0.25m^2((a + b)(a - b)) = a^2 - b^2$$

(iv) $(a + 3b)(x + 5)$

$$= a(x + 5) + 3b(x + 5)$$

$$= ax + 5a + 3bx + 15b$$

(v) $2pq(3pq - 2q^2) + 3q(3pq - 2q^2)$

$$= 6p^2q^2 - 4pq^3 + 9pq^3 - 6q^4$$

$$= 6p^2q^2 - 5pq^3 - 6q^4$$

(vi) $\frac{3}{4}a^2\left(4a^2 - \frac{8}{3}b^2\right) + 12b^2\left(a^2 - \frac{2}{3}b^2\right)$

$$= 3a^4 - 2a^2b^2 + 12a^2b^2 - 8b^4$$

$$= 3a^4 - 8b^4 + 10a^2b^2$$

#464478

Topic: Operations of Polynomials

Find the product

(i) $(5 - 2x)(3 + x)$

(ii) $(x + 7y)(7x - y)$

(iii) $(a^2 + b)(a + b^2)$

(iv) $(p^2 - q^2)(2p + q)$

Solution

$$(i) 5(3 + x) - 2x(3 + x)$$

$$= 15 + 5x - 6x - 2x$$

$$= 15 - x - 2x^2$$

$$(ii) x(7x - y) + 7y(7x - y)$$

$$= 7x^2 - xy + 49xy - 7y^2$$

$$= 7x^2 - 7y^2 + 48xy$$

$$(iii) a^2(a + b^2) + b(a + b^2)$$

$$= a^3 + a^2b^2 + ab + b^3$$

$$(iv) p^2(2p + q) - q^2(2p + q)$$

$$= 2p^3 + p^2q - 2pq^2 - q^3$$

#464485

Topic: Operations of Polynomials**Simplify**

$$(i) (x^2 - 5)(x + 5) + 25$$

$$(ii) (a^2 + 5)(b^3 + 3) + 5$$

$$(iii) (t + 8^2)(t^2 - s)$$

$$(iv) (a + b)(c - d) + (a - b)(c + d) + 2(ac + bd)$$

$$(v) (x + y)(2x + y) + (x + 2y)(x - y)$$

$$(vi) (x + y)(x^2 - xy + y^2)$$

$$(vii) (1.5x - 4y)(1.5x + 4y + 3) - 4.5x + 12y$$

$$(viii) (a + b + c)(a + b - c)$$

Solution

$$(i) x^2(x+5) - 5(x+5) + 25$$

$$= x^3 + 5x^2 - 5x$$

$$(ii) a^2b^3 + 3a^2 + 5b^3 + 15 + 5$$

$$a^2b^3 + 3a^2 + 5b^3 + 20$$

$$(iii) t(t^2 - s) + s^2(t^2 - s)$$

$$= t^3 - st + s^2t^2 - s^3$$

$$(iv) = a(c-d) + b(c-d) + a(c+d) - b(c+d) + 2(ac+bd)$$

$$ac - ad + bc - bd + ac + ad - bc - bd + 2ac + 2bd$$

$$= 4ac$$

$$(v) 2x^2 + xy + 2xy + y^2 + x^2 - xy + 2xy - 2y^2$$

$$2x^2 + x^2 + y^2 - 2y^2 + xy + 2xy - xy + 2xy$$

$$3x^2 + 4xy - y^2$$

$$(vi) x^3 - x^2y + xy^2 + x^2y - xy^2 + y^3$$

$$x^3 + y^3 + (xy^2 - xy^2) + (x^2y - x^2y)$$

$$= x^3 + y^3$$

$$(vii) 2.25x^2 + 6xy + 4.5x - 6xy - 16y^2 - 12y - 4.5x + 12y$$

$$= 2.25x^2 - 16y^2$$

$$(viii) a(a+b+c) + b(a+b-c) + c(a+b-c)$$

$$a^2 + ab - ac + ab + b^2 - bc + ac + bc - c^2$$

$$= a^2 + b^2 - c^2 + (ab + ab) + (bc - bc) + (ca - ca)$$

$$= a^2 + b^2 - c^2 + 2ab$$

#464491

Topic: Standard Identities

Use a suitable identity to get each of the following products.

(i) $(x + 3)(x + 3)$

(ii) $(2y + 5)(2y + 5)$

(iii) $(2a - 7)(2a - 7)$

(iv) $\left(3a - \frac{1}{2}\right)\left(3a - \frac{1}{2}\right)$

(v) $(1.1m - 0.4)(1.1m + 0.4)$

(vi) $(a^2 + b^2)(-a^2 + b^2)$

(vii) $(6x - 7)(6x + 7)$

(viii) $(-a + c)(-a + c)$

(ix) $\left(\frac{x}{2} + \frac{3y}{4}\right)\left(\frac{x}{2} + \frac{3y}{4}\right)$

(x) $(7a - 9b)(7a - 9b)$

Solution

$$\begin{aligned} \text{(i)} \quad & (x+3)^2 \\ & = x^2 + 6x + 9 \quad \because [(a+b)^2 = a^2 + b^2 + 2ab] \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad & (2y+5)^2 \\ & = 4y^2 + 25 + 20y \quad \because [(a+b)^2 = a^2 + b^2 + 2ab] \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad & (2a-7)^2 \\ & = 4a^2 - 28a + 49 \quad \because [(a-b)^2 = a^2 + b^2 - 2ab] \end{aligned}$$

$$\begin{aligned} \text{(iv)} \quad & 3a^2 - 2(3a)\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^2 \\ & = 9a^2 - 3a + \frac{1}{4} \quad \because [(a-b)^2 = a^2 + b^2 - 2ab] \end{aligned}$$

$$\begin{aligned} \text{(v)} \quad & (1.1m - 0.4)(1.1m + 0.4) \\ & = 1.21m^2 - 0.16 \quad \because [(a^2 - b^2) = (a+b)(a-b)] \end{aligned}$$

$$\begin{aligned} \text{(vi)} \quad & (a^2 + b^2)(-a^2 + b^2) \\ & = (b^2 + a^2)(b^2 - a^2) \\ & = b^4 - a^4 \quad \because [(a^2 - b^2) = (a+b)(a-b)] \end{aligned}$$

$$\begin{aligned} \text{(vii)} \quad & (6x-7)(6x+7) \\ & = (6x)^2 - (7)^2 \\ & = 36x^2 - 49 \quad \because [(a^2 - b^2) = (a+b)(a-b)] \end{aligned}$$

$$\begin{aligned} \text{(viii)} \quad & (-a+c)(-a+c) \\ & = (-a+c)^2 \\ & = a^2 - 2ac + c^2 \quad \because [(a-b)^2 = a^2 + b^2 - 2ab] \end{aligned}$$

$$\begin{aligned} \text{(ix)} \quad & \left(\frac{x}{2} + \frac{3y}{4}\right) \left(\frac{x}{2} + \frac{3y}{4}\right) \\ & = \left(\frac{x}{2} + \frac{3y}{4}\right)^2 \\ & = \left(\frac{x}{2}\right)^2 + 2\left(\frac{x}{2}\right)\left(\frac{3y}{4}\right) + \left(\frac{3y}{4}\right)^2 \\ & = \frac{x^2}{4} + \frac{3xy}{4} + \frac{9y^2}{16} \quad \because [(a+b)^2 = a^2 + b^2 + 2ab] \end{aligned}$$

$$\begin{aligned} \text{(x)} \quad & (7a-9b)(7a-9b) \\ & = (7a-9b)^2 \\ & = 49a^2 - 126ab + 81b^2 \quad \because [(a-b)^2 = a^2 + b^2 - 2ab] \end{aligned}$$

#464495

Topic: Standard Identities

Use the identity $(x + a)(x + b) = x^2 + (a + b)x + ab$ to find following products.

- (i) $(x + 3)(x + 7)$
- (ii) $(4x + 5)(4x + 1)$
- (iii) $(4x - 5)(4x - 1)$
- (iv) $(4x + 5)(4x - 1)$
- (v) $(2x + 5y)(2x + 3y)$
- (vi) $(2a^2 + 9)(2a^2 + 5)$
- (vii) $(xyz - 4)(xyz - 2)$

Solution

$$\begin{aligned} \text{(i)} \quad & (x + 3)(x + 7) \\ &= x^2 + (3 + 7)x + (3)(7) \\ &= x^2 + 10x + 21 \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad & (4x)^2 + (5 + 1)4x + (5)(1) \\ &= 16x^2 + 24x + 5 \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad & (4x - 5)(4x - 1) \\ &= (4x)^2 + [(-5) + (-1)]4x + (-5)(-1) \\ &= 16x^2 - 24x + 5 \end{aligned}$$

$$\begin{aligned} \text{(iv)} \quad & (4x)^2 + [(5) + (-1)]4x + (5)(-1) \\ &= 16x^2 + 16x - 5 \end{aligned}$$

$$\begin{aligned} \text{(v)} \quad & (2x)^2 + (5y + 3y)(2x) + (5y)(3y) \\ &= 4x^2 + 16xy + 15y^2 \end{aligned}$$

$$\begin{aligned} \text{(vi)} \quad & (2a^2)^2 + (9 + 5)(2a^2) + (9)(5) \\ &= 4a^2 + 28a^2 + 45 \end{aligned}$$

$$\begin{aligned} \text{(vii)} \quad & (xyz)^2 + [(-4) + (-2)](xyz) + (-4)(-2) \\ &= x^2y^2z^2 - 6xyz + 8 \end{aligned}$$

#464924

Topic: Standard Identities

Find the square of the following numbers

- (i) 32
- (ii) 35
- (iii) 86
- (iv) 93
- (v) 71
- (vi) 46

Solution

$$(i) 32^2 = (30 + 2)(30 + 2)$$

$$= 900 + 60 + 60 + 4$$

$$= 1024$$

$$(ii) 35^2 = (30 + 5)(30 + 5)$$

$$= 900 + 150 + 150 + 25$$

$$= 1225$$

$$(iii) 86^2 = (80 + 6)(80 + 6)$$

$$= 6400 + 480 + 480 + 36$$

$$= 7396$$

$$(iv) 93^2 = (90 + 3)(90 + 3)$$

$$= 8100 + 270 + 270 + 9$$

$$= 8649$$

$$(v) 71^2 = (70 + 1)(70 + 1)$$

$$= 4900 + 140 + 1$$

$$= 5041$$

$$(vi) 46^2 = (40 + 6)(40 + 6)$$

$$= 1600 + 480 + 36$$

$$= 2080 + 36$$

$$= 2116.$$

#466259

Topic: Standard Identities

Find the following squares by using the identities.

$$(i) (b - 7)^2$$

$$(ii) (xy + 3z)^2$$

$$(iii) (6x^2 - 5y)^2$$

$$(iv) \left(\frac{2}{3}m + \frac{3}{2}n\right)^2$$

$$(v) (0.4p - 0.5q)^2$$

$$(vi) (2xy + 5y)^2$$

Solution

$$\text{i)} (b - 7)^2 \\ = b^2 + 7^2 - 2(b)(7) \\ = b^2 + 49 - 14b$$

$$\therefore [(a - b)^2 = a^2 + b^2 - 2ab]$$

$$\text{ii)} (xy + 3z)^2 \\ = (xy)^2 + (3z)^2 + 2(xy)(3z) \\ = x^2y^2 + 9z^2 + 6xyz$$

$$\therefore [(a + b)^2 = a^2 + b^2 + 2ab]$$

$$\text{iii)} (6x - 5y)^2 \\ = (6x)^2 + (5y)^2 - 2(6x)(5y) \\ = 36x^2 + 25y^2 - 60xy$$

$$\therefore [(a - b)^2 = a^2 + b^2 - 2ab]$$

$$\text{iv)} \left(\frac{2}{3}m + \frac{3}{2}n\right)^2 \\ = \left(\frac{2}{3}m\right)^2 + \left(\frac{3}{2}n\right)^2 + 2\left(\frac{2}{3}m\right)\left(\frac{3}{2}n\right) \\ = \frac{4}{9}m^2 + \frac{9}{4}n^2 + 2mn$$

$$\therefore [(a + b)^2 = a^2 + b^2 + 2ab]$$

$$\text{v)} (0.4p - 0.5q)^2 \\ = (0.4p)^2 + (0.5q)^2 - 2(0.4p)(0.5q) \\ = 0.16p^2 + 0.25q^2 - 4pq$$

$$\therefore [(a - b)^2 = a^2 + b^2 - 2ab]$$

$$\text{vi)} (2xy + 5y)^2 \\ = (2xy)^2 + (5y)^2 + 2(2xy)(5y) \\ = 4x^2y^2 + 25y^2 + 20xy \\ = y^2(4x^2 + 20x + 25)$$

#466260

Topic: Standard Identities

Simplify:

- (i) $(a^2 - b^2)^2$
- (ii) $(2x + 5)^2 - (2x - 5)^2$
- (iii) $(7m - 8n)^2 + (7m + 8n)^2$
- (iv) $(4m + 5n)^2 + (4n + 5m)^2$
- (v) $(2.5p - 1.5q)^2 - (1.5p - 2.5q)^2$
- (vi) $(ab + bc)^2 - 2ab^2c$
- (vii) $(m^2 - n^2m^2 + 2m^3n^2)$

Solution

$$\text{i) } (a^2 - b^2)^2 = (a^2)^2 + (b^2)^2 - 2(a^2)(b^2) \quad \because [(a+b)^2 = a^2 + b^2 + 2ab] \\ = a^4 + b^4 - 2a^2b^2$$

$$\text{ii) } (2x+5)^2 - (2x-5)^2$$

Let $A = 2x+5$ and $B = 2x-5$

$$\therefore (2x+5)^2 - (2x-5)^2 = A^2 - B^2$$

$$= (A+B)(A-B)$$

$$= [2x+5 + (2x-5)][2x+5 - (2x-5)]$$

$$= [2x+5 + 2x-5][2x+5 - 2x+5]$$

$$= (4x)(10)$$

$$= 40x$$

$$\text{iii) } (7m-8n)^2 + (7m+8n)^2$$

$$= (7m)^2 + (8n)^2 - 2(7m)(8n) + (7m)^2 + (8n)^2 + 2(7m)(8n)$$

$$= 49m^2 + 64n^2 - 112mn + 49m^2 + 64n^2 + 112mn$$

$$= 98m^2 + 128n^2$$

$$\text{iv) } (4m+5n)^2 + (5m+4n)^2$$

$$= (4m)^2 + (5n)^2 + 2(4m)(5n) + (5m)^2 + (4n)^2 + 2(5m)(4n)$$

$$= 16m^2 + 25n^2 + 40mn + 25m^2 + 16n^2 + 40mn$$

$$= 41m^2 + 41n^2 + 80mn$$

$$\text{v) } (2.5p-1.5q)^2 - (1.5p-2.5q)^2$$

$$= (2.5p)^2 + (1.5q)^2 - 2(2.5p)(1.5q) - [(1.5p)^2 + (2.5q)^2 - 2(1.5p)(2.5q)]$$

$$= 6.25p^2 + 2.25q^2 - 7.5pq - 2.25p^2 - 6.25q^2 + 7.5pq$$

$$= 4p^2 - 4q^2$$

$$= 4(p^2 - q^2)$$

$$= 4(p-q)(p+q)$$

$$\text{vi) } (ab+bc)^2 - 2ab^2c$$

$$= (ab)^2 + (bc)^2 + 2(ab)(bc) - 2ab^2c$$

$$= a^2b^2 + b^2c^2 + 2ab^2c - 2ab^2c$$

$$= b^2(a^2 + c^2)$$

$$\text{vii) } (m^2 - n^2m)^2 + 2m^3n^2$$

$$= (m^2)^2 + (n^2m)^2 - 2(m^2)(n^2m) + 2m^3n^2$$

$$= m^4 + m^2n^4 - 2m^3n^2 + 2m^3n^2$$

$$= m^2(m^2 - n^4)$$

#466261

Topic: Standard Identities

Show that:

$$(i) (3x + 7)^2 - 84x = (3x - 7)^2$$

$$(ii) (9p - 5q)^2 + 180pq = (9p + 5q)^2$$

$$(iii) \left(\frac{4}{3}m - \frac{3}{4}n\right)^2 + 2mn = \frac{16}{9}m^2 + \frac{9}{16}n^2$$

$$(iv) (4pq + 3q)^2 - (4pq - 3q)^2 = 48pq^2$$

$$(v) (a - b)(a + b) + (b - c)(b + c) + (c - a)(c + a) = 0$$

Solution

We know that

$$(a \pm b)^2 = a^2 + b^2 \pm 2ab$$

and,

$$(a - b)(a + b) = a^2 - b^2$$

Now,

(i)

$$\begin{aligned} (3x + 7)^2 - 84x &= (3x)^2 + 7^2 + 2(3x)(7) - 84x = (3x)^2 + 7^2 + 42x - 84x \\ &= (3x)^2 + 7^2 - 42x = (3x - 7)^2 \end{aligned}$$

(ii)

$$\begin{aligned} (9p - 5q)^2 + 180pq &= (9p)^2 + (5q)^2 - 2(9p)(5q) + 180pq = (9p)^2 + (5q)^2 - 90pq + 180pq \\ &= (9p)^2 + (5q)^2 + 90pq = (9p + 5q)^2 \end{aligned}$$

(iii)

$$\begin{aligned} \left(\frac{4}{3}m - \frac{3}{4}n\right)^2 + 2mn &= \left(\frac{4}{3}m\right)^2 + \left(\frac{3}{4}n\right)^2 - 2\left(\frac{4}{3}m\right)\left(\frac{3}{4}n\right) + 2mn \\ &= \left(\frac{4}{3}m\right)^2 + \left(\frac{3}{4}n\right)^2 - 2mn + 2mn \\ &= \frac{16}{9}m^2 + \frac{9}{16}n^2 \end{aligned}$$

(iv)

$$\begin{aligned} (4pq + 3q)^2 - (4pq - 3q)^2 &= (4pq)^2 + (3q)^2 + 2(4pq)(3q) - ((4pq)^2 + (3q)^2 - 2(4pq)(3q)) \\ &= (4pq)^2 + (3q)^2 + 2(4pq)(3q) - (4pq)^2 - (3q)^2 + 2(4pq)(3q) \\ &= 24pq^2 + 24pq^2 = 48pq^2 \end{aligned}$$

(v)

$$\begin{aligned} (a - b)(a + b) + (b - c)(b + c) + (c - a)(c + a) &= a^2 - b^2 + b^2 - c^2 + c^2 - a^2 \\ &= 0 \end{aligned}$$

#466262

Topic: Special Products

Using identities, evaluate

(i) 71^2 (ii) 99^2 (iii) 102^2 (iv) 998^2

(v) 5.2^2 (vi) 297×303 (vii) 78×82

(viii) 8.9^2 (ix) 1.05×9.5

Solution

$$\begin{aligned} \text{i)} 71^2 &= (70 + 1)^2 \\ &= (70)^2 + 1^2 + 2(70)(1) \quad \because [(a + b)^2 = a^2 + b^2 + 2ab] \\ &= 4900 + 1 + 140 \end{aligned}$$

= 5041

ii) $99^2 = (100 - 1)^2$
 $= (100)^2 + 1^2 - 2(100)(1)$ $\therefore [(a - b)^2 = a^2 + b^2 - 2ab]$
 $= 10000 + 1 - 200$
 $= 9801$

iii) $102^2 = (100 + 2)^2$
 $= (100)^2 + 2^2 + 2(100)(2)$ $\therefore [(a + b)^2 = a^2 + b^2 + 2ab]$
 $= 10000 + 4 + 400$
 $= 10404$

iv) $998^2 = (1000 - 2)^2$
 $= (1000)^2 + 2^2 - 2(1000)(2)$ $\therefore [(a - b)^2 = a^2 + b^2 - 2ab]$
 $= 1000000 + 4 - 4000$
 $= 996004$

v) $5.2^2 = (5 + 0.2)^2$
 $= (5)^2 + (0.2)^2 + 2(5)(0.2)$ $\therefore [(a + b)^2 = a^2 + b^2 + 2ab]$
 $= 25 + 0.04 + 2$
 $= 27.04$

vi) $297 \times 303 = (300 - 3)(300 + 3)$
 $= (300)^2 - (3)^2$ $\therefore [(a^2 - b^2) = (a + b)(a - b)]$
 $= 90000 - 9$
 $= 89991$

vii) $78 \times 82 = (80 - 2)(80 + 2)$
 $= (80)^2 - (2)^2$ $\therefore [(a^2 - b^2) = (a + b)(a - b)]$
 $= 6400 - 4$
 $= 6396$

viii) $(8.9)^2 = (9 - 0.1)^2$
 $= (9)^2 + (0.1)^2 - 2(9)(0.1)$ $\therefore [(a - b)^2 = a^2 + b^2 - 2ab]$
 $= 81 + 0.01 - 1.8$
 $= 79.21$

vii) $1.05 \times 9.5 = \frac{105}{100} \times \frac{95}{10}$

$$= \frac{1}{1000} \times (105 \times 95)$$

$$= \frac{1}{1000} \times (100 + 5)(100 - 5)$$

$$= \frac{1}{1000} \times [(100)^2 - (5)^2]$$

$$= \frac{1}{1000} \times (10000 - 25)$$

$$= \frac{9975}{1000} = 9.975$$

#466263**Topic:** Standard IdentitiesUsing $a^2 - b^2 = (a + b)(a - b)$, find

- (i) $51^2 - 49^2$
- (ii) $1.02^2 - 0.98^2$
- (iii) $153^2 - 147^2$
- (iv) $12.1^2 - 7.9^2$

Solution

i) $51^2 - 49^2$

$= (51 - 49)(51 + 49)$

$= 2 \times 100$

$= 200$

ii) $(1.02)^2 - (0.98)^2$

$= (1.02 - 0.98)(1.02 + 0.98)$

$= 0.04 \times 2$

$= 0.08$

iii) $153^2 - 147^2$

$= (153 - 147)(153 + 147)$

$= 6 \times 300$

$= 1800$

iv) $12.1^2 - 7.9^2$

$= (12.1 - 7.9)(12.1 + 7.9)$

$= 4.2 \times 20$

$= 8.4$

We know that

$a^2 - b^2 = (a - b)(a + b)$

(i) $51^2 - 49^2 = (51 - 49)(51 + 49) = 2 \times 100 = 200$

(ii) $1.02^2 - 0.98^2 = (1.02 - 0.98)(1.02 + 0.98) = 0.04 \times 2.00 = 0.08$

(iii) $153^2 - 147^2 = (153 - 147)(153 + 147) = 6 \times 300 = 1800$

(iv) $12.1^2 - 7.9^2 = (12.1 - 7.9)(12.1 + 7.9) = 4.2 \times 20 = 84$

#466264**Topic:** Special ProductsUsing $(x + a)(x + b) = x^2 + (a + b)x + ab$, find

- (i) 103×104
- (ii) 5.1×5.2
- (iii) 103×98
- (iv) 9.7×9.8

Solution

$$\begin{aligned}\text{i)} 103 \times 104 &= (100 + 3)(100 + 4) \\ &= (100)^2 + (3 + 4)100 + (3)(4) \\ &= 10000 + 700 + 12 \\ &= 10712\end{aligned}$$

$$\begin{aligned}\text{ii)} 5.1 \times 5.2 &= (5 + 0.1)(5 + 0.2) \\ &= (5)^2 + (0.1 + 0.2)5 + (0.1)(0.2) \\ &= 25 + 1.5 + 0.02 \\ &= 26.52\end{aligned}$$

$$\begin{aligned}\text{iii)} 103 \times 98 &= (100 + 3)(100 - 2) \\ &= (100)^2 + (3 - 2)100 + (3)(-2) \\ &= 10000 + 100 - 6 \\ &= 10094\end{aligned}$$

$$\begin{aligned}\text{iv)} 9.7 \times 9.8 &= (9 + 0.7)(9 + 0.8) \\ &= (9)^2 + (0.7 + 0.8)9 + (0.7)(0.8) \\ &= 81 + 13.5 + 0.56 \\ &= 95.06\end{aligned}$$

